

DIANEAL LOW CALCIUM PERITONEAL DIALYSIS SOLUTION WITH DEXTROSE - dextrose, sodium chloride, sodium lactate, calcium chloride and magnesium chloride injection, solution
BAXTER HEALTHCARE CORPORATION

DESCRIPTION

DIANEAL Low Calcium peritoneal dialysis solutions in AMBU-FLEX containers are sterile, nonpyrogenic solutions for intraperitoneal administration only. They contain no bacteriostatic or antimicrobial agents or added buffers.

Composition, calculated osmolality, pH and ionic concentrations are shown in Table 1.

Potassium is omitted from peritoneal dialysis solutions because dialysis may be performed to correct hyperkalemia. In situations in which there is a normal serum potassium level or hypokalemia, the addition of potassium chloride (up to a concentration of 4 mEq/L) may be indicated to prevent severe hypokalemia. **Addition of potassium chloride should be made after careful evaluation of serum and total body potassium and only under the direction of a physician.** Frequent monitoring of serum electrolytes is indicated.

In some patients calcium carbonate is used as a phosphate binder. Because serum calcium levels have been observed to be elevated in these patients (Slatopolsky et al. 1986), the calcium concentration of DIANEAL Low Calcium peritoneal dialysis solutions has been reduced to 2.5 mEq/L. Serum calcium levels should be monitored and if low, the amount of oral calcium carbonate phosphate binder may be increased or peritoneal dialysis solutions containing higher calcium concentrations may be used. If serum calcium levels rise, adjustments to the dosage of the calcium carbonate phosphate binder and/or vitamin D analogs should be considered by the physician. Because average plasma magnesium levels in some chronic CAPD patients have been observed to be elevated (Nolph et al. 1981), the magnesium concentration of this formulation has been reduced to 0.5 mEq/L. Average plasma magnesium levels have not been reported for chronic IPD and CCPD patients. Serum magnesium levels should be monitored and if low, oral magnesium supplements, oral magnesium containing phosphate binders, or peritoneal dialysis solutions containing higher magnesium concentrations may be used.

Because average serum bicarbonate levels in some chronic CAPD patients (Nolph et al. 1981), some chronic IPD patients (La Greca et al. 1980), and some chronic CCPD patients (Diaz-Buxo et al. 1983), have been observed to be somewhat lower than normal values, the bicarbonate precursor (lactate) concentration of DIANEAL Low Calcium peritoneal dialysis solutions has been raised to 40 mEq/L. Serum bicarbonate levels should be monitored.

The osmolalities shown in Table 1 are calculated values. Calculated osmolality of DIANEAL Low Calcium peritoneal dialysis solution with 1.5% dextrose is 344 mOsmol/L, compared with measured values in normal human serum of 280 mOsmol/L.

The plastic container is fabricated from a specially formulated polyvinyl chloride (PL 146 Plastic). The amount of water that can permeate from inside the container into the overpouch is insufficient to affect the solution significantly. Solutions in contact with the plastic container can leach out certain of its chemical components in very small amounts within the expiration period, e.g., di-2-ethylhexyl phthalate (DEHP), up to 5 parts per million; however, the safety of the plastic has been confirmed in tests in animals according to USP biological tests for plastic containers as well as by tissue culture toxicity studies.

CLINICAL PHARMACOLOGY

Peritoneal dialysis is a procedure for removing toxic substances and metabolites normally excreted by the kidneys, and for aiding in the regulation of fluid and electrolyte balance.

The procedure is accomplished by instilling peritoneal dialysis fluid through a conduit into the peritoneal cavity. With the exception of lactate, present as a bicarbonate precursor, electrolyte concentrations in the fluid have been formulated in an attempt to normalize plasma electrolyte concentrations resulting from osmosis and diffusion across the peritoneal membrane (between the patient's plasma and the dialysis fluid). Toxic substances and metabolites, present in high concentrations in the blood, cross the peritoneal membrane into the dialyzing fluid. Dextrose in the dialyzing fluid is used to produce a solution hyperosmolar to the plasma, creating an osmotic gradient which facilitates fluid removal from the patient's plasma into the peritoneal cavity. After a period of time (dwell time), the fluid is drained from the cavity.

INDICATIONS AND USAGE

DIANEAL Low Calcium peritoneal dialysis solutions are indicated for use in chronic renal failure patients being maintained on peritoneal dialysis.

CONTRAINDICATIONS

None known.

WARNINGS

Peritoneal dialysis should be done with great care, if at all, in patients with a number of abdominal conditions including disruption of the peritoneal membrane or diaphragm by surgery or trauma, extensive adhesions, bowel distention, undiagnosed abdominal disease, abdominal wall infection, hernias or burns, fecal fistula or colostomy, tense ascites, obesity, and large polycystic kidneys (Vaamonde and Perez 1977). Other conditions include recent aortic graft replacement and severe pulmonary disease. When assessing peritoneal dialysis as the mode of therapy in such extreme situations, the benefits to the patient must be weighed against the possible complications.

An accurate fluid balance record must be kept and the weight of the patient carefully monitored to avoid over or under hydration with severe consequences including congestive heart failure, volume depletion, and shock.

Excessive use of DIANEAL Low Calcium peritoneal dialysis solution with 3.5% or 4.25% dextrose during a peritoneal dialysis treatment can result in significant removal of water from the patient.

Stable patients undergoing maintenance peritoneal dialysis should have routine periodic evaluation of blood chemistries and hematologic factors, as well as other indicators of patient status.

In some patients calcium carbonate is used as a phosphate binder. Because serum calcium levels have been observed to be elevated in these patients (Slatopolsky et al. 1986), the calcium concentration of DIANEAL Low Calcium peritoneal dialysis solutions has been reduced to 2.5 mEq/L. Serum calcium levels should be monitored and if low, the amount of oral calcium carbonate phosphate binder may be increased or peritoneal dialysis solutions containing higher calcium concentrations may be used. If serum calcium levels rise, adjustments to the dosage of the calcium carbonate phosphate binder and/or vitamin D analogs should be considered by the physician. Because average plasma magnesium levels in some chronic CAPD patients have been observed to be elevated (Nolph et al. 1981), the magnesium concentration of this formulation has been reduced to 0.5 mEq/L. Average plasma magnesium levels have not been reported for chronic IPD and CCPD patients. Serum magnesium levels should be monitored and if low, oral magnesium supplements, oral magnesium containing phosphate binders, or peritoneal dialysis solutions containing higher magnesium concentrations may be used.

Because average serum bicarbonate levels in some chronic CAPD patients (Nolph et al. 1981), some chronic IPD patients (La Greca et al. 1980), and some chronic CCPD patients (Diaz-Buxo et al. 1983), have been observed to be somewhat lower than normal values, the bicarbonate precursor (lactate) concentration of DIANEAL Low Calcium peritoneal dialysis solutions has been raised to 40 mEq/L. Serum bicarbonate levels should be monitored.

Not for use in the treatment of lactic acidosis.

Potassium is omitted from DIANEAL Low Calcium peritoneal dialysis solutions because dialysis may be performed to correct hyperkalemia. **Addition of potassium chloride should be made after careful evaluation of serum and total body potassium and only under the direction of a physician.**

The use of 5 or 6 liters of dialysis solution is not indicated in a single exchange.

Refer to manufacturer's directions accompanying drugs to obtain full information on additives.

If the resealable rubber plug on the medication port is missing or partially removed, do not use product if medication is to be added.

After the pull ring has been removed, inspect connector of solution container for fluid flow. A few drops of solution within the connector or pull ring may be present due to condensation of water resulting from the sterilization process. If a continuous stream of fluid is noted, discard solution because sterility may be impaired.

After removing overpouch, check for minute leaks by squeezing container firmly. If leaks are found, discard the solution because the sterility may be impaired.

Freezing of solution may occur at temperatures below 0°C (32°F). Allow to thaw naturally in ambient conditions and thoroughly mix contents by shaking.

PRECAUTIONS

Aseptic technique must be used throughout the procedure and at its termination in order to reduce the possibility of infection. If peritonitis occurs, the choice and dosage of antibiotics should be based upon the results of identification and sensitivity studies of the isolated organism(s) when possible. Prior to identification of the involved organism(s), broad-spectrum antibiotics may be indicated. Peritoneal dialysis solutions may be warmed in the overpouch to 37°C (98.6°F) to enhance patient comfort. However, only dry heat (for example, heating pad) should be used. Solutions should not be heated in water due to an increased risk of infection. Microwave ovens should not be used to heat solutions because there is a potential for damage to the solution container. Moreover, microwave oven heating may potentially cause overheating and/or non-uniform heating of the solution that may result in patient injury or discomfort.

Significant losses of protein, amino acids and water soluble vitamins may occur during peritoneal dialysis. Replacement therapy should be provided as necessary.

Pregnancy

Teratogenic Effects

Pregnancy Category C

Animal reproduction studies have not been conducted with DIANEAL Low Calcium peritoneal dialysis solutions. It is also not known whether DIANEAL Low Calcium peritoneal dialysis solutions can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. DIANEAL Low Calcium peritoneal dialysis solutions should be given to a pregnant woman only if clearly needed.

Do not administer unless solution is clear and seal is intact.

ADVERSE REACTIONS

Adverse reactions to peritoneal dialysis include mechanical and solution related problems as well as the results of contamination of equipment or improper technique in catheter placement. Abdominal pain, bleeding, peritonitis, subcutaneous infection around a chronic peritoneal catheter, catheter blockage, difficulty in fluid removal, and ileus are among the complications of the procedure. Solution related adverse reactions may include electrolyte and fluid imbalances, hypovolemia, hypervolemia, hypertension, hypotension, disequilibrium syndrome, and muscle cramping.

When prescribing the solution to be used for an individual patient, consideration should be given to the potential interaction between the dialysis treatment and therapy directed at other existing illnesses. For example, rapid potassium removal may create arrhythmias in cardiac patients using digitalis or similar drugs; digitalis toxicity may be masked by elevated potassium or magnesium, or by hypocalcemia. Correction of electrolytes by dialysis may precipitate signs and symptoms of digitalis excess. Conversely, toxicity may occur at suboptimal dosages of digitalis if potassium is low or calcium high. Azotemic diabetics require careful monitoring of insulin requirements during and following dialysis with dextrose containing solutions.

DOSAGE AND ADMINISTRATION

DIANEAL Low Calcium peritoneal dialysis solutions are intended for intraperitoneal administration only.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration whenever solution and container permit.

The mode of therapy (Intermittent Peritoneal Dialysis [IPD], Continuous Ambulatory Peritoneal Dialysis [CAPD], or Continuous Cyclic Peritoneal Dialysis [CCPD]), frequency of treatment, formulation, exchange volume, duration of dwell, and length of dialysis should be selected by the physician responsible for and supervising the treatment of the individual patient.

To avoid the risk of severe dehydration and hypovolemia and to minimize the loss of protein, it is advisable to select the peritoneal dialysis solution with the lowest level of osmolarity consistent with the fluid removal requirements for that exchange.

Peritoneal dialysis solutions may be warmed in the overpouch to 37°C (98.6°F) to enhance patient comfort. However, only dry heat (for example, heating pad) should be used. (See Directions for Use)

The addition of heparin to the dialysis solution may be indicated to aid in prevention of catheter blockage in patients with peritonitis, or when the solution drainage contains fibrinous or proteinaceous material (Ribot et al. 1966). 1000 to 2000 USP units of heparin per liter of solution has been recommended for adults (Furman et al. 1978). For children, 50 units of heparin per 100 mL of dialysis fluid has been recommended (Irwin et al. 1981).

Additives may be incompatible. Complete information is not available. Those additives known to be incompatible should not be used. Consult with pharmacist, if available. If, in the informed judgment of the physician, it is deemed advisable to introduce additives, use aseptic technique. Mix thoroughly when additives have been introduced. Do not store solutions containing additives.

Intermittent Peritoneal Dialysis (IPD)

For maintenance dialysis of chronic renal failure patients.

The cycle of instillation, dwell and removal of dialysis fluid is repeated sequentially over a period of hours (8 to 36 hours) as many times per week as indicated by the condition of the patient. For chronic renal failure patients, maintenance dialysis is often accomplished by periodic dialysis (3 to 5 times weekly) for shorter time periods (8 to 14 hours per session) (Mattocks and El-Bassiouni 1971).

Continuous Ambulatory Peritoneal Dialysis (CAPD) and Continuous Cyclic Peritoneal Dialysis (CCPD)

For maintenance dialysis of chronic renal failure patients.

In CAPD, typically 1.5 to 3.0 liters of dialysis solution (depending upon patient size) are instilled into the peritoneal cavity of adults and the peritoneal access device is then clamped (Kim et al. 1984; Twardowski and Janicka 1981; Twardowski and Burrows 1984). For children, 30 to 50 mL/kg body weight with a maximum of 2 liters has been recommended (Potter et al. 1981; Irwin et al. 1981). The solution remains in the cavity for dwell times of 4 to 8 hours during the day and 8 to 12 hours overnight. At the conclusion of each dwell period, the access device is opened, the solution drained and fresh solution instilled. The procedure is repeated 3 to 5 times per day, 6 to 7 days per week. Solution exchange volumes and frequency of exchanges should be individualized for adequate biochemical and fluid volume control (Moncrief et al. 1982; Twardowski et al. 1983). The majority of exchanges will utilize 1.5% or 2.5% dextrose containing peritoneal dialysis solutions, with 3.5% or 4.25% dextrose containing solutions being used when extra fluid removal is required. Patient weight is used as the indicator of the need for fluid removal (Popovich et al. 1978).

In CCPD, the patient receives 3 or 4 dialysis exchanges during the night which range from 2-1/2 to 3 hours dwell duration. Typically 1.5 to 2.0 liters of dialysis solution (depending upon patient size) are delivered each cycle by an automatic peritoneal dialysis cyclor machine. After the last outflow during the night, an additional exchange is infused by the cyclor machine into the peritoneum. The equipment is then disconnected from the patient, and the dialysate remains in the peritoneum for 14 to 15 hours during the day until the next nocturnal cycle (Diaz-Buxo et al. 1981). Combinations of 1.5% or 2.5% dextrose containing peritoneal dialysis solutions are usually used for the nighttime exchanges, while 3.5% or 4.25% dextrose containing solution is used when extra fluid removal is required such as during the daytime exchange. Patient weight is used as the indicator of the need for fluid removal (Popovich et al. 1978) so therapy should be individualized according to the patient's need for ultrafiltration.

It is recommended that adult patients being placed on peritoneal dialysis or, in the case of pediatric patients, the selected caretaker, (as well as the patient, when suitable), should be appropriately trained in a program which is under the supervision of a physician. Training materials are available from Baxter Healthcare Corporation, Deerfield, IL 60015 USA to facilitate this training.

HOW SUPPLIED

DIANEAL Low Calcium peritoneal dialysis solutions in AMBU-FLEX II and AMBU-FLEX III containers are available in nominal size flexible containers with fill volumes as indicated in Table 1.

All DIANEAL Low Calcium peritoneal dialysis solutions have overfills which are declared on container labeling.

Exposure of pharmaceutical products to heat should be minimized. Avoid excessive heat. It is recommended the product be stored at room temperature (25°C/77°F): brief exposure up to 40°C (104°F) does not adversely affect the product.

Directions for Use

Use aseptic technique.

For complete system preparation, see directions accompanying ancillary equipment.

Peritoneal dialysis solutions may be warmed in the overpouch to 37°C (98.6°F) to enhance patient comfort. However, only dry heat (for example, heating pad) should be used. Solutions should not be heated in water due to an increased risk of infection. Microwave ovens should not be used to heat solutions because there is a potential for damage to the solution container. Moreover, microwave oven heating may potentially cause overheating and/or non-uniform heating of the solution that may result in patient injury or discomfort.

To Open

Tear overpouch down side at slit and remove solution container. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. If supplemental medication is desired, follow directions below before preparing for administration. Check for minute leaks by squeezing container firmly.

To Add Medication

Additives may be incompatible.

If the resealable rubber plug on the medication port is missing or partially removed, do not use product if medication is to be added.

1. Put on mask. Clean and/or disinfect hands.
2. Prepare medication site using aseptic technique.
3. Using a syringe with a 1 inch long 19 to 25 gauge needle, puncture resealable medication port and inject medication.
4. Position container with ports up and evacuate the medication port by squeezing and tapping it.
5. Mix solution and medication thoroughly.

Preparation for Administration

1. Put on mask. Clean and/or disinfect hands.
2. Place solution container on work surface.
3. Remove pull ring from connector of the solution container. If continuous fluid flow from connector is observed, discard solution container.
4. Remove tip protector from tubing set and immediately attach to connector of the solution container.
5. Continue with therapy set-up as instructed in user manual or directions accompanying tubing sets.
6. Upon completion of therapy, discard unused portion.

REFERENCES


1. Diaz-Buxo, J.A. et al. 1981. Continuous cyclic peritoneal dialysis: a preliminary report. **Int Soc Artif Organs** 81:157-161.
2. Diaz-Buxo, J.A. et al. 1983. Observations on inadequate base buffer concentrations in peritoneal dialysis solutions. **ASAIO Abstracts** 43.
3. Furman, K.I. et al. 1978. Activity of intraperitoneal heparin during peritoneal dialysis. **Clinical Nephrology** 9:15-18.

4. Irwin, M.A. et al. 1981. Continuous ambulatory peritoneal dialysis in pediatrics. **AANNT J** 8:11-13, 44.
5. Kim, D. et al. 1984. Continuous ambulatory peritoneal dialysis with three-liter exchanges: a prospective study. **Peritoneal Dial Bull** 4:82-85.
6. La Greca, G. et al. 1980. Acid base balance on peritoneal dialysis. **Clinical Nephrology** 16(1):1-6.
7. Mattocks, A.M. and El-Bassiouni, E.A. 1971. Peritoneal dialysis: a review. **J Pharm Sci** 60:1767-1782.
8. Moncrief, J.W. et al. 1982. CAPD: Are three exchanges per day adequate? **AANNT J** 9:39-43.
9. Nolph, K.D. et al. 1981. Considerations for dialysis solution modifications. In **Peritoneal Dialysis**, eds. Robert C. Atkins et al. Chapter 25. New York: Churchill Livingstone.
10. Popovich, R.P. et al. 1978. Continuous ambulatory peritoneal dialysis. **Ann Intern Med** 8:449-456.
11. Potter, D.E. et al. 1981. Continuous ambulatory dialysis (CAPD) in children. **Trans Am Soc Artif Intern Organs** 27:64-67.
12. Ribot, S. et al. 1966. Complications of peritoneal dialysis. **Am J Med Sci** 252:505-517.
13. Slatopolsky, E. et al. 1986. Calcium carbonate as a phosphate binder in patients with chronic renal failure undergoing dialysis. **NEJM** 3:315, 157-160.
14. Twardowski, Z.J. and Janicka, L. 1981. Three exchanges with a 2.5 liter volume for continuous ambulatory peritoneal dialysis. **Kidney Int** 20:281-284.
15. Twardowski, Z.J. et al. 1983. High volume low frequency continuous ambulatory peritoneal dialysis. **Kidney Int** 23:64-70.
16. Twardowski, Z.J. and Burrows, L. 1984. Two year experience with high volume, low frequency continuous ambulatory peritoneal dialysis. **Peritoneal Dial Bull** 4:S67.
17. Vaamonde, C.A. and Perez, G.O. 1977. Peritoneal dialysis today. **Kidney** 10:31-36.

Table 1																
	Composition/100mL					OSMOLARITY (mOsmol/L) (calc)	pH	Ionic Concentration (mEq/L)					How Supplied			
	*Dextrose Hydrate USP	Sodium Chloride USP (NaCl)	Sodium Lactate USP (C ₃ H ₅ NaO ₃)	Calcium Chloride USP (CaCl ₂ •2H ₂ O)	Magnesium Chloride USP (MgCl ₂ •6H ₂ O)			Sodium	Calcium	Magnesium	Chloride	Lactate	Fill Volume (mL)	Container Size (mL)	Code	NDC
DIANEAL Low Calcium Peritoneal Dialysis Solution with 1.5% Dextrose AMBU- FLEX II Container	15 g	538 mg	448 mg	18.3 mg	5.08 mg	344	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	2000 2500 3000 5000 6000	3000 3000 3000 6000 6000	L5B4825 L5B9091 L5B9901 L5B4826 L5B9770	NDC 0941-0409-06 NDC 0941-0409-08 NDC 0941-0409-05 NDC 0941-0409-07 NDC 0941-0409-01

DIANEAL	15 g	538 mg	448 mg	18.3 mg	5.08 mg	344	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	1500	2000	5B9715	NDC
Calcium													2000	2000	5B4825	0941-0409-45
Peritoneal													2500	3000	5B9718	NDC
Dialysis													3000	3000	5B9904	0941-0409-36
Solution													5000	5000	5B4826	NDC
with													6000	6000	5B9770	0941-0409-48
1.5%																NDC
Dextrose																0941-0409-49
AMBU-																NDC
FLEX																0941-0409-27
III																NDC
Container																0941-0409-28
DIANEAL	15 g	538 mg	448 mg	18.3 mg	5.08 mg	395	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	2000	3000	L5B9727	NDC
Low													2500	3000	L5B9028	0941-0457-08
Calcium													3000	3000	L5B9902	NDC
Peritoneal													5000	6000	L5B5003	0941-0457-07
Dialysis													6000	6000	L5B9771	NDC
Solution																0941-0457-02
with																NDC
2.5%																0941-0457-05
Dextrose																NDC
AMBU-																0941-0457-01
FLEX																
II																
Container																
DIANEAL	15 g	538 mg	448 mg	18.3 mg	5.08 mg	395	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	1500	2000	5B9725	NDC
Calcium													2000	3000	5B9727	0941-0457-45
Peritoneal													2500	3000	5B9728	NDC
Dialysis													3000	3000	5B9902	0941-0457-47
Solution													5000	5000	5B5202	NDC
with													6000	6000	5B9770	0941-0457-48
2.5%																NDC
Dextrose																0941-0457-49
AMBU-																NDC
FLEX																0941-0457-25
III																NDC
Container																0941-0457-28
DIANEAL	15 g	538 mg	448 mg	18.3 mg	5.08 mg	445	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	2500	3000	5B9738	NDC
Calcium																0941-0463-48
Peritoneal																
Dialysis																
Solution																
with																
3.5%																
Dextrose																
AMBU-																
FLEX																
III																
Container																
DIANEAL	12.5 g	538 mg	448 mg	18.3 mg	5.08 mg	483	5.2 (4.0 to 6.5)	132	2.5	0.5	95	40	2000	3000	L5B9747	NDC
Low													2500	3000	L5B9028	0941-0459-08
Calcium													3000	3000	L5B9903	NDC
Peritoneal													5000	6000	L5B5003	0941-0459-07
Dialysis													6000	6000	L5B9772	NDC
Solution																0941-0459-02

with 4.25% Dextrose AMBU- FLEX II Container																NDC 0941-0459-05 NDC 0941-0459-01
DIANEAL Calcium g Peritoneal Dialysis Solution with 4.25% Dextrose AMBU- FLEX III Container	4.15 mg	538 mg	448 mg	18.3 mg	5.08 mg	483 (4.0 to 6.5)	5.2	132	2.5	0.5	95	40	1500 2000 2500 3000 5000 6000	2000 3000 3000 3000 5000 6000	5B9745 5B9746 5B9748 5B9908 5B5203 5B9742	NDC 0941-0459-45 NDC 0941-0459-47 NDC 0941-0459-48 NDC 0941-0459-49 NDC 0941-0459-25 NDC 0941-0459-28



Baxter, DIANEAL, AMBU-FLEX, and PL 146 are trademarks of
 Baxter International Inc.
Baxter Healthcare Corporation
 Deerfield, IL 60015 USA
 Printed in USA
 ©Copyright 1981, 1982, 1983, 1984, 1989, 2008 Baxter Healthcare Corporation.
 All rights reserved.
 07-19-60-956
 2009/08

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL
Container Label

5B4825

NDC 0941-0409-36



2000 mL

(APPROX 80 mL EXCESS)

Baxter

**Dianeal
Low Calcium (2.5 mEq/L)
Peritoneal Dialysis Solution
with 1.5% Dextrose**

EACH 100 mL CONTAINS 1.5 g DEXTROSE HYDROUS USP
538 mg SODIUM CHLORIDE USP 448 mg SODIUM
LACTATE 18.3 mg CALCIUM CHLORIDE USP 5.08 mg
MAGNESIUM CHLORIDE USP pH 5.2 (4.0 TO 6.5)

mEq/L SODIUM - 132 CALCIUM - 2.5 MAGNESIUM -
0.5 CHLORIDE - 95 LACTATE - 40
OSMOLARITY - 344 mOsmol/L (CALC)

STERILE NONPYROGENIC

POTASSIUM CHLORIDE TO BE ADDED ONLY
UNDER THE DIRECTION OF A PHYSICIAN

READ PACKAGE INSERT FOR FULL INFORMATION

FOR INTRAPERITONEAL ADMINISTRATION ONLY

DOSAGE AS DIRECTED BY A PHYSICIAN

CAUTIONS SQUEEZE AND INSPECT INNER BAG
WHICH MAINTAINS PRODUCT STERILITY DISCARD
IF LEAKS ARE FOUND

DO NOT USE UNLESS SOLUTION IS CLEAR

DISCARD UNUSED PORTION

Rx ONLY

STORE UNIT IN MOISTURE BARRIER OVERWRAP AT
ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO
USE

AVOID EXCESSIVE HEAT SEE INSERT

Ambu-Flex III CONTAINER PL 146 PLASTIC

BAXTER DIANEAL AMBU-FLEX III AND PL 146 ARE
TRADEMARKS OF BAXTER INTERNATIONAL INC

BAXTER HEALTHCARE CORPORATION

DEERFIELD IL 60015 USA

MADE IN USA

US PAT NO 4573980

Low Calcium 1.5% Dextrose

5B4825

NDC 0941-0409-36

2000 mL

(APPROX 80 mL EXCESS)

Baxter

Dianeal

Low Calcium (2.5 mEq/L)

Peritoneal Dialysis Solution

with 1.5% Dextrose

Low Calcium 1.5% Dextrose

EACH 100 mL CONTAINS 1.5 g DEXTROSE HYDROUS USP

538 mg SODIUM CHLORIDE USP 448 mg SODIUM
LACTATE 18.3 mg CALCIUM CHLORIDE USP 5.08 mg
MAGNESIUM CHLORIDE USP pH 5.2 (4.0 TO 6.5)
mEq/L SODIUM - 132 CALCIUM - 2.5 MAGNESIUM -
0.5 CHLORIDE - 95 LACTATE - 40
OSMOLARITY - 344 mOsmol/L (CALC)
STERILE NONPYROGENIC
POTASSIUM CHLORIDE TO BE ADDED ONLY
UNDER THE DIRECTION OF A PHYSICIAN
READ PACKAGE INSERT FOR FULL INFORMATION
FOR INTRAPERITONEAL ADMINISTRATION ONLY
DOSAGE AS DIRECTED BY A PHYSICIAN
CAUTIONS SQUEEZE AND INSPECT INNER BAG
WHICH MAINTAINS PRODUCT STERILITY DISCARD
IF LEAKS ARE FOUND
DO NOT USE UNLESS SOLUTION IS CLEAR
DISCARD UNUSED PORTION
Rx ONLY
STORE UNIT IN MOISTURE BARRIER OVERWRAP AT
ROOM TEMPERATURE (25°C/77°F) UNTIL READY TO
USE
AVOID EXCESSIVE HEAT SEE INSERT
Ambu-Flex III CONTAINER PL 146 PLASTIC
BAXTER DIANEAL AMBU-FLEX III AND PL 146 ARE
TRADEMARKS OF BAXTER INTERNATIONAL INC
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN USA
US PAT NO 4573980
Carton Label

5B4825

6-2000 ML
AMBU-FLEX™ III CONTAINERS

1.5%

DIANEAL® LOW CALCIUM 1.5% DEX
PERITONEAL DIALYSIS SOLUTION

EXP
XXXXX

SECONDARY BAR CODE

(17) YYMM00 (10) XXXXX

LOT
XXXXX

PRIMARY BAR CODE

(01) 50309410409362

5B4825

6-2000 ML

AMBU-FLEX™ III CONTAINERS

1.5%

DIANEAL® LOW CALCIUM 1.5% DEX
PERITONEAL DIALYSIS SOLUTION

EXP

XXXXX

SECONDARY BAR CODE
(17) YYMM00 (10) XXXXX
LOT
XXXXX
PRIMARY BAR CODE
(01) 50309410409362